

Clinical Guideline in Paediatric Dentistry **Stainless Steel Preformed Crowns for Primary Molars**

S A Fayle, Department of Paediatric Dentistry, Leeds Dental Institute, Clarendon Way, Leeds, LS2 9LU.

Update of guideline

S.A. Kindelan, P. Day, R. Nichol and N. Willmott, SpRs Paediatric Dentistry, Department of Paediatric Dentistry, Leeds Dental Institute, Clarendon Way, Leeds, LS2 9LU.

Since this sixth National Clinical Guideline was published in the International Journal of Paediatric Dentistry in 1999 there has been one meta-analysis¹, four literature reviews^{2,3,4,5} and one prospective clinical trial⁶ published in relation to the use of stainless steel crowns (SSCs) or preformed metal crowns (PMCs) for the restoration of deciduous molars. All papers have concluded that the failure rate for SSCs used in primary molar teeth is very low compared with plastic restorations. At the time of update of this guideline a randomised control trial (RCT) investigating the effectiveness of SSCs in managing carious primary molars using the Hall technique is underway, with one year results published.⁷

A Cochrane review pertaining to the use of preformed metal crowns for the restoration of carious primary molar teeth was published in January 2007⁸. The aim of the review was to compare clinical outcomes for primary molar teeth restored with SSCs compared to those restored with alternative filling materials or remaining untreated. Using very stringent criteria, looking for evidence from well designed RCTs, of which none could be found, the authors concluded that whilst there was a subjective impression amongst paediatric dentists that SSCs provide a more durable restoration than plastic restorative materials there is little evidence from good quality clinical trials to support this. The Cochrane report does however stress that a paucity of stringent clinical studies should not be interpreted as evidence for a lack of efficiency of the technique. Indeed whilst studies and reports published may not meet Cochrane standards it is important to emphasise that there is a large amount of useful literature advocating the use of SSCs. Butani and colleagues⁹ have described the quantity of published literature available which relate to the use of SSCs. In 2005 they found a total of 122 papers with 52 of these being outcome-related, evidence-based literature and the others reviews/ expert opinion, case reports, technique and practice guidelines.

It is essential for the clinician to use the best available evidence to support clinical practice.

This paper provides the dental practitioner with an update of the current published literature and the available evidence for the use of SSCs in the treatment of the primary molar. British Society of Paediatric Dentistry guidelines for the recommended use of SSCs in primary molar teeth are re-iterated.

Evidence underlying recommendations is scored according to the SIGN classification and guidelines should be read in this context. The process of guideline production in the UK is described in the *International Journal of Paediatric Dentistry* 1997; 7:267-268.

For this updated guideline a search of the dental literature was made electronically from MEDLINE OVID using the key words: preformed crown(s), stainless steel crown(s), (a)esthetic primary (deciduous) molar crown(s). A total of 236 abstracts were identified from 1966 to the current time. Publications pertaining to the use of stainless steel crowns in deciduous teeth

retrieved. All articles previously cited in the 1999 guideline were reviewed in addition to all papers subsequently published.

Introduction

Stainless steel (preformed) crowns are prefabricated crown forms which can be adapted to individual primary molars and cemented in place to provide a definitive restoration. The following guideline is intended to assist in the planning and provision of stainless steel crown restorations for primary molars.

1. Indications

Stainless steel crowns are the **restoration of choice** in the following situations:

- (B) 1.1 Restoration of carious primary molars where more than two surfaces are affected, or where one or two surface carious lesions are extensive.
- (B) 1.2 Following pulpotomy or pulpectomy procedures.

Stainless steel crowns may also be **indicated** in the following situations:

- (C) 1.3 Restoration of primary molars affected by localised or generalised developmental problems (e.g. enamel hypoplasia, amelogenesis imperfecta, dentinogenesis imperfecta etc.).
- (C) 1.4 Restoration of fractured primary molars.
- (C) 1.5 Restoration and protection of teeth exhibiting extensive tooth surface loss due to attrition, abrasion or erosion.
- (B) 1.6 In patients with a high caries susceptibility.
- 1.7 As an abutment for certain appliances, such as space maintainers.
- (C) 1.8 In patients where routine oral hygiene measures are impaired e.g. patients with special needs, and breakdown of intra-coronal restorations is likely.
- (C) 1.9 In patients undergoing restorative care under general anaesthesia if two or more surfaces are involved.
- (C) 2.0 In patients with infra-occluded primary molars to maintain mesiodistal space.

Stainless steel crowns are **contra-indicated**

- if the primary molar is close to exfoliation with more than half the roots resorbed
- in a patient with a known nickel allergy or sensitivity (see footnote at end of paper)

2. Clinical procedure

- 2.1 Appropriate local analgesia should be obtained and the tooth should be isolated, preferably with rubber dam.
 - 2.2 Caries removal and appropriate pulp treatment (i.e. indirect pulp capping, pulpotomy or pulpectomy) should be completed if necessary. Some clinicians advocate preparation of the tooth for the crown prior to finalising caries removal and/ or pulpotomy treatment.
- (C)
- 2.3 Appropriate tooth preparation should be carried out, which should include sufficient occlusal reduction to avoid significant occlusal prematurity, and approximal reduction to allow the crown to be seated beyond the maximum bulbosity of the crown. Occlusal reduction should follow the contours of the tooth. The preparation should finish with a smooth feather edge cervically with no step or shoulder. The preparation should be rounded off with no sharp line angles. Where a primary molar has no adjacent tooth either mesially or distally it is still important to carry out approximal reduction to avoid producing an excessive marginal overhang. This is particularly important on the distal surface of second primary molars where such overhangs can impede the eruption of the first permanent molar. Buccal and lingual preparation is not always necessary and may be detrimental to retention.
 - 2.4 A crown should be selected that is a tight snap fit. Choosing the correct size is assisted by measuring the mesio-distal dimension of the tooth, or contralateral tooth, with dividers or a graduated periodontal probe.
- (B)
- 2.5 Stainless steel crowns produced by several different manufacturers are available in the United Kingdom. The degree of adjustment necessary to achieve a satisfactory fit is dependant upon the make of crown used. SSCs crowns from 3M™ ESPE™ are anatomically trimmed and contoured cervically and in many instances require little or no modification. Other types of SSC have little or no cervical contouring and hence routinely require modification.
 - 2.51 If the crown is excessively long, the crown margin may impede complete seating, in which case crown length may be adjusted by trimming with crown shears and re-smoothing and polishing the edges with an abrasive stone. Although it has been customary to recommend trimming of crowns where gingival blanching occurs, there is no evidence that this practice reduces post cementation complications. Manufacturers recommend the SSC finishes about 1mm below the gingival margin.

- 2.52 Over trimming of the crown margin should be avoided, as this may affect retention if it results in reduced adaptation of the crown margin into undercut areas. It is essential that the margins of the crown are well adapted into undercut areas, which is usually achieved by crimping of the crown edges.
- 2.53 Special attention should be given to adaptation of the distal margin on second primary molars where the permanent molar is unerupted. An uncorrected distal overhang may result in impaction of the first permanent molar. Care should be taken not to cause iatrogenic damage to adjacent teeth or unerupted teeth.
- (C) 2.6 Frequently, reduction in the mesio-distal dimension of the crown will be necessary, especially where mesial drift (often due to caries) has resulted in loss of arch length. Moderate reduction in mesio-distal dimension can be achieved by flattening of the mesial and distal contact areas of the crown with Adam's pattern pliers. Where mesial drift has occurred in the lower arch it may be possible to use a SSC form for the contralateral upper tooth (e.g. ULE crown form for LRE) as these SSC forms have a shorter mesiodistal dimension. Other forms of modification, including vertically slicing one aspect of the crown and spot-welding additional segments of stainless steel band to increase the perimeter or extend the length have been described, but their efficacy remains largely untested.
- (C) 2.7 Excessive occlusal interference should be avoided (greater than 1.0-1.5 mm), but a slightly premature or high occlusal contact up to about 1.0mm is normally well tolerated in children, who appear to have considerable capacity for dentoalveolar compensation, with the occlusion adapting to any prematurity within a few weeks.
- (C) 2.8 The crown should be cemented with a luting cement. Glass ionomer, zinc polycarboxylate and zinc phosphate cements are all suggested by manufacturers, although fluoride-leaching cements may have added benefits. There is, however, some evidence suggesting that the specific choice of cement does not significantly affect retention, the most important retentive components being derived from correct contouring and crimping of the crown.
- (B) 2.9 Careful attention should be paid to removal of excess cement. This can usually be effectively achieved by running a pointed instrument around the margins of the cemented crown and by passing knotted dental floss bucco-lingually through the contact areas prior to the cement setting. Excess cement has been shown to be detrimental to gingival health.

3. Other Considerations

- 3.1 Stainless steel crowns may be aesthetically improved by placement of composite resin in a buccal window cut into the labial face of the crown post-fitting. Alternatively aesthetic crowns with prefabricated tooth coloured buccal and occlusal facings are available from specialist suppliers.

- (B) 3.2 When cementing orthodontic bands to stainless steel crowns roughening of the internal surface of the band and external surface of the crown prior to cementation has been shown to improve retention.

Explanatory Notes

- 1.0 Stainless steel crowns are widely recognised as the most effective and durable restoration for primary molars. There have been several retrospective studies examining the longevity of stainless steel crowns in comparison with amalgam restorations.^{10, 11, 12, 13} All have shown stainless steel crowns to have markedly superior longevity when compared with multi-surface amalgam restorations. Retrospective data suggests that stainless steel crowns similarly out-perform glass ionomer cements and composite restorations.^{14, 15, 16} Two retrospective studies have involved data collected from patients who had undergone comprehensive care of the primary dentition under general anaesthesia.^{15, 16} These authors strongly recommend the use of SSCs for the restoration of carious primary molars under general anaesthesia.
- Eriksson¹⁷ reported a non-randomised clinical trial comparing SSCs to contralateral teeth which were either restored or sound, however, it is unclear whether this trial was retrospective or prospective. When Randall¹ used this study in her meta-analysis but excluded the sound contralateral control teeth SSCs had a significantly higher success rate (78.8%) than amalgams (21.4%) over the 7-year period of the study. Roberts and Sherriff¹⁸ provided a prospective report on the survival of amalgam and SSC molar restorations placed in specialist paediatric dental practice over 10 years. The true failure rate for primary molar restorations was 4.1% for class I amalgams and 11.6% for class II amalgams, while only 1.9% of SSCs failed. The authors gave a 5-year estimated survival rate of 92% for SSCs and 67% for minimal class II cavities restored with amalgam. A more recent prospective study carried out by Roberts and colleagues⁶ over a 7-year period gave a 97% success rate for SSCs under the conditions of a specialist paediatric practice. Resin modified glass ionomers were equally as successful as SSCs when used in small class I and minimal class II cavities. Although this study was prospective it was not a randomised control trial as the treatment provided was dictated by the clinical status of the tooth, such that extensive caries was restored with a SSC whilst minimal cavities were restored with resin modified glass ionomer cement. One retrospective study has investigated the longevity of restorations placed in primary molars within an NHS practice¹⁹. Wong and Day screened 361 records of three dentists working in NHS practice, randomly analysing one restoration per patient. They found evidence that SSCs performed better than alternative restorations within the conditions of NHS practice.
- In a systematic review and meta-analysis of clinical studies comparing SSCs with amalgam restorations Randall¹ demonstrated the clinical effectiveness of SSCs over amalgam restorations for the treatment of large carious lesions in primary molars. Several of the studies cited allocated SSC treatments to restore large, multisurface carious lesions, whilst amalgam was reserved for the smaller lesions. Thus it can be seen that SSCs outperformed amalgam restorations even when placed in more demanding circumstances, adding a negative bias to the outcome of the crowned teeth and strengthening the evidence for the clinical performance of SSCs.

Despite this favourable outcome, they are seldom used in general dental practice.²⁰ Many dentists avoid the use of SSCs due to lack of clinical experience and the belief that provision of a SSC is a complex procedure, whilst in reality it is often simpler and more cost-effective treatment modality than a class II restoration. Given the longevity of SSCs,¹⁸ a primary molar treated in this way is unlikely to need any further treatment until it exfoliates naturally. It has been suggested that postgraduate training packages, and in particular hands-on training courses, may be the most effective way to encourage dental practitioners to re-evaluate the restorative techniques they undertake and may increase the use of SSCs for the restoration of primary molar teeth.²¹ Currently a randomised control clinical trial (RCT) is being carried out in Scotland to investigate the success of cementing SSCs over unprepared carious primary molars (Hall Technique) as a therapeutic option for the pre-co-operative child.^{7,22, 23} This has followed a retrospective audit of this technique which found a success rate of 67.6% after 5 years.²⁴ The results of the RCT are awaited before the technique can be recommended.

- 1.2 Retrospective studies have shown the success rate of formocresol pulpotomies to be greater for teeth restored with SSCs compared to those restored with amalgam,²⁵ composite^{26,27} or IRM.²⁸ In addition indirect pulp therapy in primary molars has been shown to be more successful where the definitive restoration was a SSC.²⁹

- 2.3 A study by Rector and co-workers³⁰ failed to demonstrate that the type of tooth preparation affected retention. In an earlier study,³¹ however, preparations maintaining the greatest surface area of buccal and lingual tooth structure were shown to be most retentive. This suggests that buccal and lingual reduction does not have any advantage with regard to retention and may even be detrimental. Studies have failed to show any increase in supra-gingival plaque accumulation associated with stainless steel crowns^{32, 33, 34} except in instances where crowns with defective margins have been placed, or where excess cement has been retained.^{35, 36} Several studies have investigated gingival health in association with stainless steel crown restorations. Two have suggested higher levels of gingivitis around teeth restored with stainless steel crowns.^{36, 37} In both these studies, however, no direct comparison was made with unrestored matched control teeth. In two studies where matched control teeth were used no difference in the level of gingivitis around stainless steel crowns was demonstrated.^{33, 34} The relationship between gingivitis and marginal defects, such as poor marginal adaptation and incomplete removal of excess cement, has been clearly demonstrated by several workers.^{32, 33, 34, 35, 36} Careful adaptation of crown margins before fitting is thus essential and the incidence of post-fitting gingivitis may be reduced by careful polishing of the crown margin.³⁸ The presence of a well-adapted SSC on a second deciduous molar does not affect the periodontal health of the neighbouring first permanent molar.³⁹ These clinical findings are confirmed by a more recent retrospective study evaluating clinically and radiographically the effect on gingival and bone structures of cemented SSCs in a sample of 177 children followed up from one to 38 month.⁴⁰ No deleterious effect on gingivae or bone occurred in the presence of good oral hygiene.

- 2.52 It has been demonstrated that close adaptation of the metal margins of the crown in the undercut areas significantly enhances retention.³⁰
The impaction of first permanent molars beneath over-hanging distal margins on poorly adapted stainless steel crowns has been reported.⁴¹ Careful attention should thus be paid to adaptation of the distal margin on second primary molars where the permanent molar is unerupted.
- 2.7 Slightly premature or high occlusal contact seems to be well tolerated in the primary and early mixed dentition and clinically appears to be compensated for within a few weeks. It is probable that there is an adaptation of the dento-alveolar complex to this occlusal interference in the growing child.
- 2.8 An *in vitro* study comparing stainless steel crown retention with polycarboxylate and glass ionomer cement failed to demonstrate any difference,⁴² and in an extensive study which demonstrated a 92% five-year survival¹⁸ the majority of crowns were cemented using a reinforced zinc oxide cement. Choice of cement would therefore appear to be non-critical. Since these guidelines were published a further *in vivo* study has failed to show a significant difference in retention of SSCs cemented with glass ionomer, zinc phosphate and zinc polycarboxylate, although there was only an eight month follow-up.⁴³
- 2.9 Where excess cement has been retained, stainless steel crowns have been shown to be associated with an increased degree of plaque accumulation.^{35,36} The relationship between gingivitis and marginal defects, such as poor marginal adaptation and incomplete removal of excess cement, has been clearly demonstrated by several workers.^{32,33,34,35,36}
- 3.1 Some parents or patients may complain about the appearance of SSCs. Aesthetic improvement of the appearance of stainless steel crowns by placement of composite resin in a buccal window cut into the labial face of the crown after cementing, has been reported in a case report of a modified SSC followed to exfoliation 23 months later, without evidence of deterioration.⁴¹
Alternatives to this technique are prefabricated tooth coloured crowns supplied by various manufacturers (e.g NuSmile® primary crowns). These require significantly increased space and consequently more preparation due to their greater bulk. With these tooth coloured crowns, manufacturers' instructions advise avoiding crimping of the crown which may make the facing susceptible to fracture. Consequently the tooth is prepared to fit the most appropriate crown. Prefabricated crowns with aesthetic facings have been shown to be prone to fracture *in vitro*.⁴⁵ A pilot study comparing 11 aesthetic crowns with 11 conventional SSC found the aesthetic crowns were bulkier, more expensive, resulted in poorer gingival health and lacked a natural appearance.⁴⁶ After a 4 year follow-up all the aesthetic crowns showed chipping of the facing.⁴⁷ More recently Yilmaz⁴⁸ compared the clinical success of SSCs made aesthetic by open facing with those which had aesthetic veneers. 18 open-faced and 15 veneered crowns were placed and followed up for 18 months. When loss of more than a third of the facing was recorded as a failure, open-faced crowns showed a 95% success, while the veneered crowns showed a success of 80%. The literature supporting the use of the aesthetic modified or prefabricated crown for the deciduous molar remains modest,

with small numbers of patients included in studies. Cost and increased chairside time are unlikely to lead to widespread use of these restorations. It is however useful for the clinician to be aware of this option and include it in the armamentarium of treatment options for occasional use. The patient should be warned about some gradual deterioration in appearance over time.

- 3.2 Orthodontic band retention on stainless steel crowns has been shown to be poorer than on unrestored teeth. Roughening of the internal surface of the band and external surface of the crown prior to cementation has been shown to improve retention strength to a level comparable with those obtained on unrestored permanent molar and premolar teeth.⁴⁹

Summary

The literature discussing SSCs from 1975 to the present day comes largely from retrospective clinical data, involving differing populations of patients, different makes of SSC, varying clinical conditions, luting cements and a multitude of operators. Although the quality of some of the literature may not meet modern day expectations, it is still valuable data which lends support to the longevity and cost-effectiveness of a restorative technique that has been available since the 1940s.

The recently published Cochrane review⁸ identified no randomised control trials which compared removal of dental caries followed by placement of a SSC with restoration using a plastic material or indeed no treatment but a conclusion was reached that there is some evidence from clinical studies of poor to medium quality that SSCs may last longer than fillings for carious deciduous teeth. All reported study results have been in agreement that SSCs outperform plastic restorations when used to restore multisurface carious lesions in primary molar teeth.

The Cochrane review called for well controlled clinical trials to properly test the efficacy of the SSC. It may, however, be difficult to attain ethical approval to test a restorative technique that has shown extremely favourable success rates in all studies cited. It would be very difficult to justify restoring a deciduous molar requiring a large multisurface restoration with an alternative material, or leaving it untreated in order to compare this to primary molars restored with SSCs.

The SSC should continue to be used to restore the deciduous primary molar.

(Footnote: The ESPE SSC consists of a chromium-nickel steel of surgical quality. Although nickel should not be released in significant amounts under normal clinical conditions ESPE (Service Centre, Germany) recommend that they are not used in patients with nickel allergies except after consultation with an allergologist or dermatologist.)

References

1. Randall RC, Vrijhoef MMA, Wilson NHF. Efficacy of preformed metal crowns vs. amalgam restorations in primary molars: a systematic review. *Journal of the American Dental Association* 2000; 131:337-343.
2. Randall RC. Preformed metal crowns for primary and permanent molar teeth: review of the literature. *Pediatric Dentistry* 2002; 24(5):489-500.
3. Seale SN. The use of stainless steel crowns. *Pediatric Dentistry* 2002; 24(5):501-505.
4. Hickel R, Kaaden C, Paschos E et al. Longevity of occlusally-stressed restorations in the posterior primary teeth. *American Journal of Dentistry* 2005; 18: 198-211.
5. Attari N, Roberts JF. Restoration of primary teeth with crowns: a systematic review of the literature. *European Archives of Paediatric Dentistry* 2006; 7(2):58-63.
6. Roberts JF, Attari N, Sherriff M. The survival of resin modified glass ionomer and stainless steel crown restorations in primary molars, placed in a specialist dental practice. *British Dental Journal* 2005; 198(7):427-431
7. Innes N, Evans DJP, Stirrups DR. The Hall technique: a randomised controlled clinical trial of a novel method of managing carious primary molars in general dental practice. One year results. *International Journal of Pediatric Dentistry* 2005; 15, supplement 2; 48; 14
8. Innes NPT, Ricketts DNJ, Evans DJP. Preformed metal crowns for decayed primary molar teeth. *Cochrane Database of Systematic Reviews* 2007, Issue 1. Art. No.:CD005512. DOI:10.1002/14651858.CD00512.pub2.
9. Butani YB, Levy SM, Nowak AJ, Kanellis MJ, Heller K, Hartz AJ, Dawson DV, Watkins CA. Overview of the evidence for clinical interventions in Pediatric Dentistry. *Pediatric Dentistry* 2005; 27; 6-11.
10. Braff MH. A comparison between stainless steel crowns and multisurface restorations in primary molars. *Journal of Dentistry for Children* 1975; 42: 474-8
11. Dawson LT, Simon JR, Taylor PP. The use of amalgam and stainless steel crown restorations for primary molars. *Journal of Dentistry for Children* 1981; 48: 420-2
12. Messer LB, Levering NJ. The durability of primary molar restorations: Observations and predictions of success of stainless steel crowns. *Pediatric Dentistry* 1988; 10(2):81-85.
13. Einwag J, Dunninger P. Stainless steel crown versus multisurface crown restorations: an 8 year longitudinal clinical study. *Quintessence International* 1996; 27: 321-3.
14. Papatthanasiou AG, Curzon ME, Fairpo CG. The influence of restorative material on the survival rate of restorations in primary molars. *Pediatric Dentistry* 1994; 16: 282-288
15. O'Sullivan EA, Curzon MEJ. The efficacy of comprehensive dental care for children under general anaesthesia. *British Dental Journal* 1991;171:56-58.
16. Tate AR., Ng MW., Needleman HL, Acs G. Failure rates of restorative procedures following dental rehabilitation under general anaesthesia. *Pediatric Dentistry* 2002; 24:1;69-71.
17. Eriksson AL, Paunio P, Isotupa K. Restoration of deciduous molars with ion-crowns: retention and subsequent treatment. *Proclamation of Finnish Dental Society* 1988;84(2):95-99.
18. Roberts JF, Sheriff M. The fate and survival of amalgams and preformed crown molar restorations placed in specialist paediatric dental practice. *British Dental Journal* 1990; 169: 237-44.
19. Wong FS, Day SJ. An investigation of factors influencing the longevity of restorations in primary molars. *Journal of the International Association of Dentistry for Children*. 1990 Jul;20(1):11-16.

20. Threfall AG, Pilkington L, Milsom KM et al. General dental practitioners' views on the use of stainless steel crowns to restore primary molars. *British Dental Journal* 2005;199(7);453-455.
21. Maggs-Rapport FL, Treasure ET, Chadwick BL. Community dental officer's use and knowledge of restorative techniques for primary molars: an audit of two Trusts in Wales. *International Journal Paediatric Dentistry* 2000;10:133-139.
22. Innes NPT, Evans DJP, Ricketts DNJ. Preformed metal crowns for decayed primary molar teeth. (Protocol) The Cochrane Database of Systematic Reviews 2005, Issue 4. Art. No.: CD005512. DOI: 10.1002/14651858.CD005512.
23. Evans DJP, Innes NPT, Stirrups DR. The Hall technique: a randomised controlled clinical trial of a novel method of managing carious primary molars in general dental practice. Acceptability of the technique. *International Journal of Paediatric Dentistry* 2005; 15, supplement 2; 50; 14
24. Innes NPT, Stirrups DR, Evans DJP, Hall N, Leggate M. A novel technique using preformed metal crowns for managing carious primary molars in general practice-A retrospective analysis. *British Dental Journal* 2006;200:451-454.
25. Holan G, Fuks A, Keltz N. Success rate of formocresol pulpotomy in primary molars restored with SSC vs amalgam. *Pediatric Dentistry* 2002;24(3);212-216.
26. Gruythusen RJM, Weerheijm KL. Calcium hydroxide pulpotomy with a light-cured cavity-sealing material after 2 years. *Journal of Dentistry for Children*. 1987;64:251-253.
27. Guelmann M, McIlwain MF., Primosch RE. Radiographic assessment of primary molar pulpotomies restored with resin-based materials. *Pediatric dentistry* 2005;27:24-27.
28. Farooq NS, Coll JA, Kuwabara A, et al. Success rates of formocresol pulpotomy and indirect pulp therapy in the treatment of deep dentinal caries in primary teeth. *Pediatric Dentistry* 2000; 22: 278-286.
29. Al-Zayer MA, Straffon LH., Feigal RJ, et al. Indirect pulp treatment of primary posterior teeth: a retrospective study. *Pediatric Dentistry* 2003;25:1;29-36.
30. Rector JA, Mitchell RJ, Spedding RH. The influence of tooth preparation and crown manipulation on the mechanical retention of stainless steel crowns. *Journal of Dentistry for Children* 1985; 52: 422-7.
31. Savide NL, Caputo AA, Luke LS. The effect of tooth preparation on the retention of stainless steel crowns. *Journal of Dentistry for Children* 1980; 46: 385-9.
32. Webber DL. Gingival health following placement of stainless steel crowns. *Journal of Dentistry for Children* 1974; 41: 186-9.
33. Machen DE et al. The effect of stainless steel crowns on marginal gingival tissue. *Journal of Dental Research* 1980; 59 (spec issue): Abst 239.
34. Durr DP, Ashrafi MH, Duncan WK. A study of plaque accumulation and gingival health surrounding stainless steel crowns. *Journal of Dentistry for Children* 1982; 49: 343-6.
35. Myers DR. A clinical study of the response of the gingival tissue surrounding Stainless Steel Crowns. *Journal of Dentistry for Children* 1975; 42: 281-4.
36. Goto G, Imanishi T, Machida Y. Clinical evaluation of preformed crown for deciduous molars. *Bulletin of Tokyo Dental College* 1970; 11: 169-76.
37. Henderson HZ. Evaluation of the preformed stainless steel crown. *Journal of Dentistry for Children* 1973; 40: 353-8.
38. Martens LC, Dermaut LR. The marginal polishing of Ion Nichro crowns: a preliminary report. *Journal of Dentistry for Children* 1983, 50: 417-21.

39. Guelmann M, Matsson L, Bimstein E. Periodontal health at first permanent molars adjacent to primary molar stainless steel crowns. *Journal of Clinical Periodontology* 1988; 15: 531-533.
40. Sharaf AA, Farsi NM. A clinical and radiographic evaluation of stainless steel crowns for primary molars. *Journal of Dentistry* 2004; 32; 27-33.
41. Croll TP, McKay MS, Castaldi CR. Impaction of permanent posterior teeth by overextended stainless steel crown margins. *Journal of Pedodontics* 1981; 5: 240-4.
42. Noffsinger DP, Jedrychowski JR, Caputo AA. Effects of polycarboxylate and glass ionomer cements on stainless steel crown retention. *Pediatric Dentistry* 1983; 5: 68-71.
43. Khinda VI, Grewal, N. Retentive efficacy of glass ionomer, zinc phosphate and zinc polycarboxylate luting cements in preformed stainless steel crowns: a comparative clinical study. *Journal of the Indian Society of Pedodontics and Preventive Dentistry* 2002; 20(2):41-6.
44. Roberts JF. The open-face stainless steel crown for primary molars. *Journal of Dentistry for Children* 1983; 50: 262-3.
45. Baker LH, Moon P, Murray JJ. Retention of esthetic veneers on primary stainless steel crowns. *Journal of Dentistry for Children* 1996; 63, 185-189.
46. Fuk AB, Ram D, Eidelman E. Clinical performance of esthetic posterior crowns in primary molars a pilot study. *Pediatric Dentistry* 1999;21: 445-448.
47. Ram D, Fuk AB, Eidelman E. Long term clinical performance of esthetic primary molar crowns. *Pediatric Dentistry* 2003;25:582-584.
48. Yilmaz Y, Kocogullari M. Clinical evaluation of two different methods of stainless steel esthetic crowns. *Journal of Dentistry for Children* 2004; 71: 212-214.
49. Beemer RL, Ferracane JL, Howard HE. Orthodontic band retention on primary molar stainless steel crowns. *Pediatric Dentistry* 1993; 15: 408-13.

Executive Summary

Clinical Guideline in Paediatric Dentistry

Stainless Steel Preformed Crowns for Primary Molars

S A Fayle, Department of Paediatric Dentistry, Leeds Dental Institute, Clarendon Way, Leeds, LS2 9LU. 0113 2336194

Update of guideline

S.A. Kindelan, P. Day, R. Nichol and N. Willmott, SpRs Paediatric Dentistry, Department of Paediatric Dentistry, Leeds Dental Institute, Clarendon Way, Leeds, LS2 9LU.

Since this sixth National Clinical Guideline was published in the International Journal of Paediatric Dentistry in 1999 there has been one meta-analysis¹, four literature reviews and one prospective clinical trial published in relation to the use of stainless steel crowns (SSCs) or preformed metal crowns (PMCs) for the restoration of deciduous molars^{2,3,4,5,6}. All these publications have expressed agreement that the failure rate for SSCs used in primary molar teeth is very low compared with plastic restorations. At the time of update of this guideline a randomised control trial (RCT) investigating the effectiveness of SSCs in managing carious primary molars using the Hall technique is underway and one year results have been published.⁷ A recent Cochrane review,⁸ although unable to find any evidence to unequivocally prove the superiority of restoring the primary molar with a SSC in comparison to other restorative materials did stress that a lack of stringent clinical studies should not be interpreted as evidence for a lack of efficiency of SSCs. Given the wide acceptance of the technique by specialists in the field of paediatric dentistry for restoration of the primary molar it would be difficult to design an ethical RCT to compare restoration of the broken down deciduous molar with a SSC to a plastic restorative material or to no treatment.

It should be emphasised that it is important for the clinician to use the best available evidence available to support clinical practice. All published literature to date cites good success rates for restoration of the primary molar with SSCs and longevity appears to surpass that for restoration with alternative plastic filling materials.

SSCs should continue to be recommended for restoration of the deciduous molar tooth in the following circumstances:

- Restoration of carious primary molars where more than two surfaces are affected, or where one or two surface carious lesions are extensive.
- Following pulpotomy or pulpectomy procedures.
- Restoration of primary molars affected by localised or generalised developmental problems (e.g. enamel hypoplasia, amelogenesis imperfecta, dentinogenesis imperfecta etc.).
- Restoration of fractured primary molars
- Restoration and protection of teeth exhibiting extensive tooth surface loss due to attrition, abrasion or erosion.
- In patients with a high caries susceptibility.
- As an abutment for certain appliances, such as space maintainers.
- In patients where routine oral hygiene measures are impaired e.g. patients with special needs, and breakdown of intra-coronal restorations is likely.
- In patients undergoing restorative care under general anaesthesia if two or more surfaces are involved.

1. Randall RC, Vrijhoef MMA, Wilson NHF. Efficacy of preformed metal crowns vs. amalgam restorations in primary molars: a systematic review. *Journal of the American Dental Association* 2000; 131:337-343.
2. Randall RC. Preformed metal crowns for primary and permanent molar teeth: review of the literature. *Pediatric Dentistry* 2002; 24(5):489-500.
3. Seale SN. The use of stainless steel crowns. *Pediatric Dentistry* 2002; 24(5):501-505.
4. Hickel R, Kaaden C, Paschos E et al. Longevity of occlusally-stressed restorations in the posterior primary teeth. *American Journal of Dentistry* 2005; 18: 198-211.
5. Attari N, Roberts JF. Restoration of primary teeth with crowns: a systematic review of the literature. *European Archives of Paediatric Dentistry* 2006; 7(2):58-63.
6. Roberts JF, Attari N, Sherriff M. The survival of resin modified glass ionomer and stainless steel crown restorations in primary molars, placed in a specialist dental practice. *British Dental Journal* 2005; 198(7):427-431
7. Innes N, Evans DJP, Stirrups DR. The Hall technique: a randomised controlled clinical trial of a novel method of managing carious primary molars in general dental practice. One year results. *International Journal of Pediatric Dentistry* 2005; 15, supplement 2; 48; 14
8. Innes NPT, Ricketts DNJ, Evans DJP. Preformed metal crowns for decayed primary molar teeth. Cochrane Database of Systematic Reviews 2007, Issue 1. Art. No.:CD005512. DOI:10.1002/14651858.CD00512.pub2.